

Version 29th August 1996

Collective Kondo effect in the Anderson–Hubbard lattice

P. Fazekas and K. Itai

Research Institute for Solid State Physics

P.O. Box 49, Budapest 114, H–1525 Hungary

Abstract. The periodic Anderson model is extended by switching on a Hubbard U for the conduction electrons. We use the Gutzwiller variational method to study the nearly integral valent limit. The lattice Kondo energy contains the U -dependent chemical potential of the Hubbard subsystem in the exponent, and the correlation-induced band narrowing in the prefactor. Both effects tend to suppress the Kondo scale, which can be understood to result from the blocking of hybridization. At half-filling, we find a Brinkman–Rice-type transition from a Kondo insulator to a Mott insulator.

Presented at the SCES '96 (Zürich, 1996)

To appear in Physica B

Recently, there has been a great deal of interest in many-band models which describe hybridization between different subsystems of strongly correlated electrons. Schork and Fulde introduced the problem of the Kondo-compensation of a magnetic impurity by a correlated band [1]. The corresponding periodic model can be envisaged either as an array of localized spins coupled to a Hubbard band (the Kondo–Hubbard lattice [2]), or as a periodic system of Anderson impurities hybridized with a Hubbard band (the Anderson–Hubbard lattice [3]). A particular version of the latter model was suggested for the remarkable correlated semiconductor FeSi [4]. We note that a closely related class of two-band models is the subject of intense investigation in connection with the resurgent interest in metallic ferromagnetism [5, 6].

In contrast to their single-impurity counterparts, the Anderson–Hubbard (AH) and Kondo–Hubbard (KH) lattice models are not equivalent [3, 7]. In fact, they tend to lead to different predictions as to the effect of the conduction band Hubbard U on the Kondo energy. A strong enhancement of the Kondo scale is found for the KH model, both for the single impurity [8] and the lattice [2]. As for AH models, there is at least one contribution which indicates the reduction of the Kondo scale with increasing U for the impurity [1, 9], and there is an argument for a reduced scale in the lattice [3]. The latter finding may be understood as the consequence of the blocking of hybridization processes.

We consider the periodic Anderson–Hubbard model which describes an array of strongly correlated f -sites hybridized with a moderately strongly interacting d -band:

$$\begin{aligned} \mathcal{H} = & \sum_{\mathbf{k},\sigma} \epsilon_d(\mathbf{k}) d_{\mathbf{k}\sigma}^\dagger d_{\mathbf{k}\sigma} + \epsilon_f \sum_{\mathbf{j},\sigma} \hat{n}_{\mathbf{j}\sigma}^f + U_f \sum_{\mathbf{j}} \hat{n}_{\mathbf{j}\uparrow}^f \hat{n}_{\mathbf{j}\downarrow}^f \\ & + U_d \sum_{\mathbf{j}} \hat{n}_{\mathbf{j}\uparrow}^d \hat{n}_{\mathbf{j}\downarrow}^d - v \sum_{\mathbf{j},\sigma} (f_{\mathbf{j}\sigma}^\dagger d_{\mathbf{j}\sigma} + d_{\mathbf{j}\sigma}^\dagger f_{\mathbf{j}\sigma}) \end{aligned} \quad (1)$$

where $\hat{n}_{\mathbf{j}\sigma}^f = f_{\mathbf{j}\sigma}^\dagger f_{\mathbf{j}\sigma}$, etc., the $\mathbf{k}$ are wave vectors, and the $\mathbf{j}$ are site indices. The d -bandwidth is W . In what follows, we take the strongly asymmetric Anderson model with $U_f \rightarrow \infty$ and the f -level $\epsilon_f < 0$ sufficiently deep-lying so that we are in the Kondo limit: $1 - n_f \ll 1$ where the f -valence is defined as $n_f = \langle \sum_{\sigma} \hat{n}_{\mathbf{j}\sigma}^f \rangle$. The total electron density (per site, for one spin) is $n = \langle \sum_{\mathbf{j},\sigma} \hat{n}_{\mathbf{j}\sigma}^f + \sum_{\mathbf{j},\sigma} \hat{n}_{\mathbf{j}\sigma}^d \rangle / 2L$, where L is the number of lattice sites. We will assume $1/2 \leq n \leq 1$, so that there are enough electrons to fill at least the f -levels, and the d -band filling is variable up to half-filling.

We use the Gutzwiller variational method, generalizing a previous treatment of the periodic Anderson model [10]. We postulate a Gutzwiller-projected hybridized band ground state

$$|\Psi\rangle = \hat{P}_G^d \cdot \hat{P}_G^f \cdot \prod_{\mathbf{k}} \prod_{\sigma} [u_{\mathbf{k}} f_{\mathbf{k}\sigma}^\dagger + v_{\mathbf{k}} d_{\mathbf{k}\sigma}^\dagger] |0\rangle \quad (2)$$

where the mixing amplitudes $u_{\mathbf{k}}/v_{\mathbf{k}}$ are treated as independent variational parameters. The Gutzwiller projector for the d -electrons is

$$\hat{P}_G^d = \prod_{\mathbf{g}} [1 - (1 - \eta) \hat{n}_{\mathbf{g}\uparrow}^d \hat{n}_{\mathbf{g}\downarrow}^d] \quad (3)$$

where the variational parameter η is controlled by U_d . For the f -electrons the full Gutzwiller projection is taken $\hat{P}_G^f = \prod_{\mathbf{g}} [1 - \hat{n}_{\mathbf{g}\uparrow}^f \hat{n}_{\mathbf{g}\downarrow}^f]$. Here we consider only non-magnetic solutions corresponding to a mass-enhanced metal at $n < 1$, or a renormalized-hybridization-gap insulator at $n = 1$.

Omitting the details of the derivation (which relies on the Gutzwiller approximation), we quote our essential results. The expression for the optimized total energy density is

$$\begin{aligned} \mathcal{E} = & \epsilon_f - n_d^0(1 - n_d^0)q_d + U_d\nu_d^0 \\ & - Wn_d^0 q_d^0 \cdot \exp \left\{ -\frac{\mu_0(U_d) - \epsilon_f}{4(v^2/W)} \right\}. \end{aligned} \quad (4)$$

The first line gives the energies of decoupled f -, and d -electrons. The coupling between the two subsystems is described by the last term which we identify as the Kondo energy of the Anderson–Hubbard model. The same Kondo scale can be identified in the deviation of the valence from 1

$$1 - n_f = \frac{n_d^0 q_d^0}{4(v/W)^2} \cdot \exp \left\{ -\frac{\mu_0(U_d) - \epsilon_f}{4(v^2/W)} \right\}. \quad (5)$$

In the above equations, $n_d^0 = n - 1/2$ is the $v = 0$ value of the conduction electron density (per spin). ν_d^0 is double occupation, q_d^0 is the q -factor (the renormalization factor of the hopping amplitude) taken with n_d^0 , and $\mu_0(U_d)$ is the chemical potential of the Hubbard band calculated (in the Gutzwiller approximation) for band filling n_d^0 .

Eqs. (5) and (4) are as yet formal results: they express the solution in terms of the optimized parameters of the Hubbard subsystem. There is no

closed-from result away from half-filling; the results of small- U_d , and of large- U_d , expansions are given in [3]. Here we discuss first the general features, and then the special case of half-filling.

The conduction band U_d appears in the characteristic Kondo scale in two ways: through the prefactor and the exponent. The prefactor describes the correlation-induced narrowing of the d -band. The exponential factor can be written as $\exp(-W/J_{\text{eff}})$ where we introduced the effective Kondo coupling

$$J_{\text{eff}} = \frac{4v^2}{\mu_0(U_d) - \epsilon_f} . \quad (6)$$

It differs from the Anderson lattice result [10, 11] inasmuch as $\mu_0(0) = (n_d^0 - 1/2)W$ is replaced by the U_d -dependent chemical potential $\mu_0(U_d)$. Comparing to the single impurity result [1], we call attention to the extra factor of 2 in the numerator: the ‘lattice enhancement of the Kondo effect’ [11] works also in the present model.

Due to the combined effect of the prefactor and the exponent, the lattice Kondo energy is a decreasing function of U_d . To make this finding plausible, one can argue that increasing U_d suppresses charge fluctuations, and therefore blocks the hybridization processes which give rise to the Kondo coupling, and thus to the Kondo effect. The blocking of hybridization is carried to an extreme in the case of exact half-filling, as we discuss below:

According to a usual Luttinger’s theorem argument, the half-filled case $n = 1$ belongs to a non-magnetic insulator. The well-known $q_d = 1 -$

$(U_d/2W)^2$ holds right up to $U_d = 2W$ where the d -band undergoes a Brinkman–Rice transition [12]. However, in contrast to the usual one-band Brinkman–Rice transition which is a metal–insulator transition, here we have to do with an insulator–insulator transition. For $U_d < 2W$, the system is a new kind of Kondo insulator with a renormalized hybridization gap [3]. For $U_d > 2W$, the system is integral valent ($n_f = 1$), the Kondo effect is completely quenched, and we should speak about a Mott insulator. This is certainly what we expect within the Gutzwiller approximation. Though the well-known criticism concerning the complete suppression of polarity fluctuations can be raised here, we note that up to the Brinkman–Rice transition, the Gutzwiller result is in a surprisingly good agreement with the predicted high-dimensional behaviour of the Hubbard model [13]. The variationally derived lattice enhancement of the Kondo scale is at least qualitatively confirmed by a similar calculation for the Anderson lattice [14]. Therefore we expect that our results which combine Hubbard and Anderson lattice features, have a similar justification for $U_d < 2W$. For $U_d > 2W$, a treatment based on replacing the Hubbard band with a t – J [15] or a Heisenberg model [16] may be more appropriate.

Our trial state is the lattice generalization of a lowest-order Varma–Yafet state [17] which is a potential source of shortcomings. Work on the impurity case is in a more advanced state and indicates that processes which enhance the Kondo scale, may predominate [8, 9]. It has been argued [9] that to

capture this effect variationally, one should consider states which are higher in the Varma–Yafet hierarchy. If a similar statement proves true for the lattice, our work served to demonstrate the limitations of the Ansatz (2). However, we believe that the lattice case may be different, for the following reason: By postulating a large Fermi surface [18], states from the entire Brillouin zone are used, and therefore (2) does contain contributions from electron–hole pairs with respect to the small (conduction electron) Fermi surface.

Acknowledgement. P. F. gratefully acknowledges financial support by the Hungarian National Science Research Foundation grant OTKA T-014201.

References

- [1] T. Schork and P. Fulde, Phys. Rev. B 50 (1994) 1345.
- [2] N. Shibata et al., Phys. Rev. B 53 (1996) R8828.
- [3] K. Itai and P. Fazekas, Phys. Rev. B 54 (1996) R752.
- [4] C. Fu and S. Doniach, Phys. Rev. B 51 (1995) 17439.
- [5] Hal Tasaki, Phys. Rev. Lett. 75 (1995) 4678.
- [6] K. Penc, H. Shiba, F. Mila, and T. Tsukagoshi: Phys. Rev. B 54 (1996) 4056.

- [7] Y.M. Li, Phys. Rev. B 52 (1995) 6979.
- [8] G. Khaliullin and P. Fulde, Phys. Rev. B 52 (1995) 9514.
- [9] T. Schork: Phys. Rev. B 53 (1996) 5626.
- [10] P. Fazekas and B.H. Brandow, Phys. Scr. 36 (1987) 809.
- [11] T.M. Rice and K. Ueda, Phys. Rev. B 36 (1986) 6420.
- [12] W.F. Brinkman and T.M. Rice, Phys. Rev. B 2 (1970) 4302.
- [13] G. Moeller et al., Phys. Rev. Lett. 74 (1995) 2082.
- [14] M. Jarrell, Phys. Rev. B 51 (1995) 7429.
- [15] S. Moukouri et al., Phys. Rev. B 53 (1996) R488.
- [16] J. Igarashi et al., Phys. Rev. B 51 (1995) 5814; *ibid* 52 (1995) 15966.
- [17] C.M. Varma and Y. Yafet, Phys. Rev. B 13 (1976) 2950.
- [18] H. Shiba and P. Fazekas, Progr. Theor. Phys. Suppl. **101**, 403 (1990).